

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.



Defects in the images may include (but are not limited to):

- BLACK BORDERS
 - TEXT CUT OFF AT TOP, BOTTOM OR SIDES
 - FADED TEXT
 - ILLEGIBLE TEXT
-
- SKEWED/SLANTED IMAGES
 - COLORED PHOTOS
 - BLACK OR VERY BLACK AND WHITE DARK PHOTOS
 - GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

08XNCCNG ***** FIRM 10032
00000029091660AS-12/01/95 N-2895 14
COLUMBIA UNIV
HEALTH SCIENCES LIB
701 N 168TH ST
NEW YORK NY 10032-2704





164
Risky science



214
Pretty productive

NEWS

- Physicists Create New State of Matter ☒ 152
- Plasma Physics: Go Back to Basics, Says NRC Panel 153
- More Than One Way to Fuse a Plasma 154
- Hughes Tosses Lifeline to 90 Eastern European Scientists 155
- Space Science: House Panel Targets Centers, Cassini 156
- Imanishi-Kari Case: Baltimore Defends Paper at Center of Misconduct Case 157
- Heavy Weather Ahead for Clinical Research 158
- New Faculty May Lose Family Tuition Help 158
- Marine Center Is Lightning Rod in Dispute Over Restoration 159
- Taking the Pulse of the Sun in Records of the Solar Wind 160
- How the T_H2 Response Is Marshaled ☒ 161
- ## SPECIAL NEWS REPORT
- Epidemiology Faces Its Limits 164
- Sizing Up the Cancer Risks 165
- Press Coverage: Leaving Out the Big Picture 166

PERSPECTIVES

- An Intimate Gathering of Bosons ☒ 18
- K. Burnett
- Methyl Chloroform and the Atmosphere ☒ 18
- A. R. Ravishankara and D. L. Albritton
- CD1: Presenting Unusual Antigens to Unusual T Lymphocytes ☒ 18
- A. Bendelac

ARTICLE

- Atmospheric Trends and Lifetime of CH_3CCl_3 and Global OH Concentrations ☒ 18
- R. G. Prinn, R. F. Weiss, B. R. Miller, J. Huang, F. N. Alyea, D. M. Cunnold, P. J. Fraser, D. E. Hartley, P. G. Simmonds

RESEARCH ARTICLE

- Protein Folding Intermediates: Native-State Hydrogen Exchange 19
- Y. Bai, T. R. Sosnick, L. Mayne, S. W. Englander

REPORTS

- Observation of Bose-Einstein Condensation in a Dilute Atomic Vapor ☒ 19
- M. H. Anderson, J. R. Ensher, M. R. Matthews, C. E. Wieman, E. A. Cornell

THIS WEEK IN SCIENCE

- EDITORIAL 141
- The Politics of Science 143
- J. H. Gibbons
- LETTERS 145
- Succeeding Generations: C. T. Hill, J. Maddox, M. Heylin, E. Rubinstein, S. Mitton • Science and Political Reality: R. S. Walker • China's "Missing" Girls: T. O. Cheng, S. Tullapurkar • New Light on Free Electron Lasers: A. Gover, W. van Amersfoort, W. B. Colson, 146

DEPARTMENTS

- K. Mima, R. Warren • Applied Research in South Africa: G. F. R. Ellis • Correction: M. E. Gurney
- SCIENCESCOPE 151
- RANDOM SAMPLES 163
- BOOK REVIEWS 253
- Powering Apollo, reviewed by A. Roland • Vignettes • Reprints of Books • Books Received
- PRODUCTS & MATERIALS 255

Board of Reviewing Editors

Frederick W. Alt
Don L. Anderson
Michael Ashburner
Stephen J. Benkovic
David E. Bloom
Piet Borst
Henry R. Bourne
Michael S. Brown
James J. Bull
Kathryn Calame

C. Thomas Caskey
Dennis W. Choi
John M. Coffin
F. Fleming Crim
Paul J. Crutzen
James E. Dahlberg
Robert Desimone
Paul T. Englund
Richard G. Fairbanks
Douglas T. Fearon

Harry A. Fazzard
Klaus Friedrich
Theodore H. Geballe
Roger I. M. Glass
Stephen P. Goff
Peter N. Goodfellow
Corey S. Goodman
Ira Herskowitz
Tomas Hökfelt
Eric F. Johnson

Stephen M. Kosslyn
Michael LaBarbera
Nicole Le Douarin
Charles S. Levings III
Alexander Levitzki
Harvey F. Lodish
Richard Losick
Reinhard Lührmann
Diane Mathis
Anthony R. Means

Shigetada Nakanishi
Roger A. Nicoll
Stuart L. Pimm
Yeshayau Pocker
Dennis A. Powers
Ralph S. Quatrano
V. Ramanathan
Douglas C. Roes
T. M. Rice
David C. Rubie

Erkki Ruoslahti
Gottfried Schatz
Jozsef Schell
Ronald H. Schwartz
Terrence J. Sejnowski
Ellen Solomon
Thomas A. Steitz
Michael P. Stryker
Robert T. N. Tjian
Emil R. Unanue

Geerat J. Vermeij
Bert Vogelstein
Arthur Weiss
Zena Werb
George M. Whitesides
Owen N. Witte
William A. Wulf

color image of the velocity distribution in a cloud of sodium atoms that have formed a Bose-Einstein condensate. Color indicates the density of atoms having a velocity specified by the two horizontal axes. The density blue and white spire is an image of low-

energy atoms that have condensed into a single quantum state. The average speed of the atoms in the spire is about 0.5 millimeter per second. See page 198 and the related News story on page 152 and the Perspective on page 182. [Image: M. R. Matthews]



Strategy for a Convergent Synthesis of Linked Glycopeptides on a Solid Support 202
Roberge, X. Beebe, S. J. Danishefsky

Motor-Enhanced Electron Transfer: Cytochrome c as a Redox-Inert Probe of Protein Complexes 204
Zhou, J. M. Nocek, M. L. DeVan, B. M. Hoffman

Experimental Studies and Theoretical Calculations for the $H + D_2 \rightarrow HD + D$ Reaction 207
Knieper, K. Seekamp-Rahn, J. Borkowski, Wrede, K. H. Welge, F. J. Aoi, L. Bañares, D'Mello, V. J. Herrero, V. Sáez Rábanos, R. E. tt

Tree Records Covering the Last Glaciation 210
Hewitt and M. Bender

Isotopic Changes in the Partial Pressure of Carbon Dioxide in Coral Reef Water 214
Meybeck, A. Suzuki, H. Saito

Formation of Diamond in Mantle Melt at High Pressure 216
Suzuki, E. Ohtani, T. Kato

Related Reductions in Human Memory Due to Impaired Encoding 218
Grady, A. R. McIntosh, B. Horwitz, J. M. Ogilvie, L. G. Ungerleider, M. J. Mentis, P. Pietrini, J. Schapiro, J. V. Haxby

Activation of Human Leukocyte Tyrosine Kinases Through Protein-Coupled Receptors 221
Knaus, S. Morris, H.-J. Dong, J. Chernoff, J. Bokoch

CD1 Binding and Presentation by CD1 223
Castafio, S. Tangri, J. E. W. Miller, H. R. Ombre, M. R. Jackson, W. D. Huse, M. J. Berg, P. A. Peterson

CD1-Restricted T Cell Recognition of Microbial Lipoglycan Antigens 227
P. A. Sieling, D. Chatterjee, S. A. Porcelli, T. I. Prigozy, R. J. Mazzaccaro, T. Soriano, B. R. Bloom, M. B. Brenner, M. Kronenberg, P. J. Brennan, R. L. Modlin

Targeted Disruption of Mouse EGF Receptor: Effect of Genetic Background on Mutant Phenotype 230
D. W. Threadgill, A. A. Dlugosz, L. A. Hansen, T. Tennenbaum, U. Lichti, D. Yee, C. LaMantia, T. Mouton, K. Herrup, R. C. Harris, J. A. Barnard, S. H. Yuspa, R. J. Coffey, T. Magnuson

Strain-Dependent Epithelial Defects in Mice Lacking the EGF Receptor 234
M. Sibilio and E. F. Wagner

Requirement of the Yeast *RTH1* 5' to 3' Exonuclease for the Stability of Simple Repetitive DNA 238
R. E. Johnson, G. K. Kovvali, L. Prakash, S. Prakash


High Concentrations of Toxaphene in Fishes from a Subarctic Lake 240
K. A. Kidd, D. W. Schindler, D. C. G. Muir, W. L. Lockhart, R. H. Hesselein

A Role in B Cell Activation for CD22 and the Protein Tyrosine Phosphatase SHP 242
G. M. Doody, L. B. Justement, C. C. Delibrias, R. J. Matthews, J. Lin, M. L. Thomas, D. T. Fearon

Lack of Interferon γ Receptor β Chain and the Prevention of Interferon γ Signaling in T_H1 Cells 245
A. Pernis, S. Gupta, K. J. Gollob, E. Garfein, R. L. Coffman, C. Schindler, P. Rothman

TECHNICAL COMMENTS

Seasonal Precipitation Timing and Ice Core Records 247
C. D. Charles, D. Rind, J. Jouzel, R. D. Koster, R. G. Fairbanks



185, 223, & 227
CD1 in skin

Board of Directors

I. Ayala
President
n
hwell
it
anco
it-elect

ester Jr.
win
lovacek

Anna C. Roosevelt
Alan Schriesheim
Jean E. Taylor
Chang-Lin Tien
Nancy S. Wexler

William T. Golden
Treasurer
Richard S. Nicholson
Executive Officer

■ SCIENCE (ISSN 0036-8075) is published weekly on Friday, except the last week in December, by the American Association for the Advancement of Science, 1333 H Street, NW, Washington, DC 20005. Second-class postage (publication No. 484460) paid at Washington, DC, and additional mailing offices. Copyright © 1995 by the American Association for the Advancement of Science. The title SCIENCE is a registered trademark of the AAAS. Domestic individual membership and subscription (51 issues): \$97 (\$50 allocated to subscription). Domestic institutional subscription (51 issues): \$228. Foreign postage extra: Mexico, Caribbean (surface mail) \$53; other countries (air assist delivery) \$93. First class, airmail, student and emeritus rates on request. Canadian rates with GST available upon request. GST #1254 88122. Printed in the U.S.A.

Change of address: allow 6 weeks, giving old and new addresses and 11-digit account number. Postmaster: Send change of address to Science, P.O. Box 2033, Marion, OH 43305-2033. Single copy sales: \$7.00 per issue prepaid includes surface postage; bulk rates on request. Authorization to photocopy material for internal or personal use under circumstances not falling within the fair use provisions of the Copyright Act is granted by AAAS to libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that \$3.00 per article is paid directly to CCC, 27 Congress Street, Salem, MA 01970. The identification code for Science is 0036-8075/93 \$3.00. Science is indexed in the Reader's Guide to Periodical Literature and in several specialized indexes.

A Strategy for a Convergent Synthesis of N-Linked Glycopeptides on a Solid Support

Jacques Y. Roberge, Xenia Beebe, Samuel J. Danishefsky

Oligosaccharides and glycopeptides are of considerable importance in molecular biology and pharmacology. However, their synthesis is complicated by the large number of different linking sites between each saccharide unit, the need for stereochemical control, the chemical sensitivity of the glycopeptide bonds, and the need to harmonize diverse protecting groups. Here, an efficient solid-phase synthesis of three N-linked glycopeptides based on glycal assembly is presented. The peptide domain can be extended while the ensemble remains bound to the polymer. The glycopeptides synthesized here are among the largest N-linked glycopeptides ever accessed by either solution- or solid-phase synthesis.

The surge of interest in glycoproteins (1-3) arises from heightened awareness of their importance in diverse biochemical processes, including cell growth regulation, binding of pathogens to cells (4), intercellular communication, and metastasis (5). Glycoproteins serve as cell differentiation markers and assist in protein folding and transport, possibly by providing protection against proteolysis (6). Improved isolation techniques and structural elucidation methods (7) have revealed high levels of microheterogeneity in naturally produced glycoproteins (8). Single eukaryotic cell lines often produce many glycoforms of any given protein sequence. For instance, erythropoietin, a clinically useful red blood cell stimulant against anemia, is glycosylated by more than 13 known types of oligosaccharide chains when expressed in Chinese hamster ovary cells (9). The efficacy of erythropoietin is heavily dependent on the type and extent of glycosylation (10).

Elucidation of the biological relevance of particular glycoprotein oligosaccharide chains benefits from isolation of pure entities. Glycoprotein heterogeneity renders this process particularly labor intensive. Some relief may be in sight in that particular cell lines can be selected to produce more homogeneous glycoproteins for structure-activity relation studies (11). However, the problem of isolation from natural sources remains daunting.


Fortunately, receptors normally recognize only a small fraction of a given macromolecular glycoconjugate. Consequently, synthesis of smaller but well-defined putative glycopeptide ligands could emerge as competitive with isolation as a source of

critical structural information (9). Important progress in glycopeptide synthesis pioneered by Kunz and others allowed synthetic access to homogenous target systems both in solution and in the solid phase (2, 12-15). Cohen-Anisfeld and Lansbury have recently reported a highly convergent solution-based coupling of selected already-available saccharides with peptides (13). In our method, the terminal glycal on the solid phase is linked with a peptide domain to generate an asparagine-linked N-acetylglucosamine construct; the whole ensemble is then retrieved and deblocked. This procedure allows one to fashion the carbohydrate domain of choice and also benefits from the


advantages associated with solid-phase synthesis in the critical carbohydrate-peptide coupling step.

The most successful approach to solid phase N-linked glycopeptide synthesis in current practice involves construction of peptide segment bearing an amine at the terminal residue on a solid support. The amine is then deprotected and coupled with either an appropriate oligosaccharide or small glycopeptide. Cleavage from the solid support and deprotection yields the desired glycopeptide (12).

The method we propose is illustrated in Scheme 1. An oligosaccharide terminating in a glycal is constructed on the solid support (see structure 1), which can be an extended linear structure or can contain branching as desired (16, 17). Through chemistry described below (Scheme 2), 1 is converted to the solid phase-bound 2, bearing a terminal 2-N-acetyl-1 β -aminoglucosamine residue (GlcNAc). A peptide is readily assembled through standard solution-phase peptide synthesis methodology or by a solid-phase assembly-retrieval sequence. Coupling of 2 with a suitable aspartic acid-containing peptide affords solid phase-bound glycopeptide 3. Retrieval and full deprotection affords the desired N-linked glycopeptide. In addition, deprotection of the carboxy terminus and addition of a peptide with a free amino terminus allows for elongation of the peptide chain while the glycopep-



Scheme 1. N-Linked glycopeptide synthesis through the use of polymer-bound glycals.



Scheme 2. Preparation of polymer-bound 2-N-acetyl-1 β -amino glucosylamine by the azasulfonamidation reaction. P = saccharide protecting group; "I⁺" = iodonium bis(collidine) perchlorate; "N₃⁻" = tetrabutylammonium azide (Bu₄NN₃); Anth = anthracene; [a] acylation with acetic anhydride (Ac₂O) and 4-N,N-dimethylaminopyridine (DMAP); [b] reduction with 1,3-propanedithiol and N,N-diisopropyl-N'-ethylamine (i-Pr₂NEt).

J. Y. Roberge and X. Beebe, Laboratory for Bioorganic Chemistry, Sloan-Kettering Institute for Cancer Research, 1275 York Avenue, Box 106, New York, NY 10021, USA.

S. J. Danishefsky, Laboratory for Bioorganic Chemistry, Sloan-Kettering Institute for Cancer Research, 1275 York Avenue, Box 106, New York, NY 10021, and Department of Chemistry, Columbia University, New York, NY 10027, USA.

de 3 is still bound to the solid support.

A simple sequence was devised to convert 1 to 2 (Scheme 2). The use of the anthracenesulfonamide (18) in the azasulfonamidation sequence (19) was crucial for the addition step (4), the azide-induced rearrangement (5), and the presentation of the solid phase-bound GlcNAc residue bearing a 1 β -amino function (2). Two examples of this design for glycopeptide syn-

thesis are illustrated in Scheme 3, showing the relative simplicity of protecting-group requirements and the high order of convergence of the approach. They result, after deprotection, in 22 or 23.

The principal advantage in using the anthracenesulfonamide is that the nitrogen-sulfur linkage can be cleaved by a variety of mild methods (18). For instance, we developed the use of thiophenol or 1,3-


propanedithiol and Hunig's base for the removal of the anthracenesulfonyl group. These protocols are compatible with the solid-supported synthesis. Also, anthracenesulfonamide itself is more soluble than benzenesulfonamide in tetrahydrofuran (THF), which is a good swelling solvent for the polymer-supported work. Thus, the use of the anthracene-based agent results in a more efficient and complete iodosulfonamidation reaction.

In solution-phase coupling of carbohydrates and peptides, the process of separating the unreacted components and by-products is not a trivial matter. Purification is greatly simplified by conducting the coupling reaction on the solid support. Most of the excess peptide is recovered from the washing by chromatography. Small amounts are lost when the activated aspartic residue cyclizes to an aspartimide. In practice, protected trisaccharide pentapeptide 21 was obtained in 37% overall yield on the basis of the initial loading of galactal carbonate. The average yield for each of the 10 steps of the sequence to 21 was 91%. Chromatography on a short column of reverse-phase silica (octadecylsilane) was sufficient to obtain this compound in pure form. This purification capability arises from the previously described "self-policing" feature of the solid-phase glycal assembly method, which avoids deletions through destruction of uncoupled donors before the next coupling cycle (17).


The glycopeptides retrieved from the support were deprotected as shown and the fully deblocked glycopeptides 22 and 23 were obtained in 61% and 48% overall yields from 20 and 21, respectively. Structural characterization of the glycopeptides by nuclear magnetic resonance spectroscopy showed the β configuration of all the anomeric linkages. The structures were further corroborated by mass spectroscopy.

The presence of orthogonal protecting groups on the carboxyl and amino termini of the peptide provides the opportunity to extend the peptide chain while the ensemble is bound to the solid support. Alternatively, after removal from the support, the freed peptide terminus may provide a functionality for linking to a carrier molecule to generate other glycoconjugates (20). Scheme 4 shows how the peptide portion of the glycopeptide was extended while still bound to the polymer support. Solid phase-bound trisaccharide pentapeptide 24 was assembled and the carboxyl terminus deprotected to give the acid, 25. Polymer-bound 25 was then coupled to tripeptide 29 with a free amino terminus to give glycopeptide 26. Retrieval from the solid support afforded trisaccharide octapeptide 27 in an 18% overall yield from polymer-bound galactal carbonate.

Regarding the future of glycopeptide



Scheme 3. Solid-phase synthesis of *N*-linked glycopeptides. All = allyl; Troc = 2,2,2-trichloroethoxy-carbonyl; Cbz = benzyloxycarbonyl; Bn = benzyl. [a] Dimethyldioxirane, THF (23), 7 or 8, zinc chloride, and THF; [b] Ac₂O, collidine, DMAP, and THF; [c] tetrabutylammoniumfluoride, acetic acid, and THF at 0°C; [d] iodonium bis(collidine) perchlorate (24) and AnthSO₂NH₂ (9 → 11); [e] Bu₄NN₃ and THF (11 → 2); [f] Ac₂O, DMAP, and THF; [g] thiophenol, *i*-Pr₂NEt, and THF; [h] 1,3-propanedithiol, *i*-Pr₂NEt (14), and THF (13 → 14); [i] 1-isobutoxy-2-isobutoxycarbonyl-1,2-dihydroquinoline (IIDQ), CH₂Cl₂, and 16 or 7 (14 + 16 → 18 and 14 + 17 → 19); [j] HF-pyridine, anisole (25), and THF (18 → 20 and 19 → 21); [k] d:PPh₃, dimethylbarbituric acid, and THF (26); [l] Zn, acetic acid, and methanol (26); [m] H₂, Pd(OAc)₂, and methanol (27); [n] KCN and methanol. Yields 22 (61% from 20) or 23 (48% from 21) (28).



Scheme 4. Extension of the peptide portion of the glycopeptide on the solid support. PMB = *p*-methoxybenzyl. [a] **28**, IIDQ, and CH₂Cl₂ (**14** → **24**); [b] Pd(PPh₃)₄, dimethylbarbituric acid, and THF; [c] **29**, IIDQ, and CH₂Cl₂; [d] HF-pyridine, anisole, and CH₂Cl₂ (**26** → **27**).

synthesis, one should also consider the method of Wong, which involves enzymatically mediated elaboration of a solid phase-bound glycosylated peptide construct using glycosyl transferases to append unprotected nucleoside phosphate-activated monosaccharides in the elongation phase (21). By this method, retrieval from the solid support is mediated by protease action. The relief from protecting groups in enzymatically mediated chemistry can be a substantial advantage. Both Wong's method and our method allow the use of unnatural amino acids and non-amino acids. The method described herein is, in principle, totally general in that it does not require the existence of the transferases and the availability of nucleoside-activated hexoses. It can also accommodate the inclusion of unnatural (artificial) sugars in the scheme. Such building blocks are available from the Lewis acid-catalyzed diene-aldehyde cyclocondensation reaction (22). All workable approaches, whether purely chemical or chemoenzymatic, are complementary for reaching the common goal of carefully designed, fully synthetic glycopeptides.

REFERENCES AND NOTES

- M. J. McPherson, P. Quirke, G. R. Taylor, Eds., *PCR: A Practical Approach* (IRL Press, Oxford, UK, 1991); G. M. Blackburn and M. J. Gait, Eds., *Nucleic Acids in Chemistry and Biology* (IRL Press, Oxford, UK, 1990); A. M. Bray, A. G. Jhingran, R. M. Valerio, N. J. Maaji, *J. Org. Chem.* **59**, 2197 (1994); G. Jung and A. G. Beck-Sickinger, *Angew. Chem. Int. Ed. Engl.* **31**, 367 (1992); M. A. Gallop, R. W. Barrett, W. J. Dower, S. P. A. Fodor, E. M. Gordon, *J. Med. Chem.* **37**, 1233 (1994); H. P. Nestler, P. A. Bartlett, W. C. Still, *J. Org. Chem.* **59**, 4723 (1994).
- M. Meldal, *Curr. Opin. Struct. Biol.* **4**, 710 (1994).
- T. Feizi and D. Bundle, *ibid.*, p. 673.
- O. P. Bahl, in *Glycoconjugates: Composition, Structure, and Function*, H. J. Allen and E. C. Kisailus, Eds. (Dekker, New York, 1992), p. 1-12.
- A. Kobata, *Acc. Chem. Res.* **26**, 319 (1993).
- G. Opdenakker, P. M. Rudd, C. P. Ponting, R. A.

Dwek, *FASEB J.* **7**, 1330 (1993).

- A. Dell and K.-H. Khoo, *Curr. Opin. Struct. Biol.* **3**, 687 (1993).
- R. A. Dwek, C. J. Edge, D. J. Harvey, M. R. Wormald, R. B. Parekh, *Annu. Rev. Biochem.* **62**, 65 (1993).
- Y. C. Lee and R. T. Lee, Eds., *Neoglycoconjugates: Preparation and Applications* (Academic Press, London, 1994).
- E. Watson, A. Bhide, H. van Halbeek, *Glycobiology* **4**, 227 (1994).
- M. A. Lehman and Z. Yucheng, U.S. Patent 5 272 070 A (1993).
- M. Meldal, in (9), p. 145; S. J. Danishefsky and J. Y. Roberge, in *Glycopeptides and Related Compounds: Chemical Synthesis, Analysis and Applications*, D. G. Large and C. D. Warren, Eds. (Dekker, New York, in press).

- S. T. Cohen-Anisfeld and P. T. Lansbury Jr., *J. Am. Chem. Soc.* **115**, 10531 (1993).
- S. T. Anisfeld and P. T. Lansbury Jr., *J. Org. Chem.* **55**, 5560 (1990).
- D. Vetter, D. Turnely, S. K. Singh, M. A. Gallop, *Angew. Chem. Int. Ed. Engl.* **34**, 60 (1995).
- J. T. Randolph, K. F. McClure, S. J. Danishefsky, *J. Am. Chem. Soc.*, in press; J. T. Randolph and S. J. Danishefsky, *Angew. Chem. Int. Ed. Engl.* **33**, 1470 (1994); S. J. Danishefsky, J. T. Randolph, J. Y. Roberge, K. F. McClure, R. B. Ruggeri, in *The Schering Lecture Series* (Schering, Germany, 1995); *Polym. Prepr. Am. Chem. Soc. Div. Polym. Chem.* **35**, 977 (1994).
- S. J. Danishefsky, K. F. McClure, J. T. Randolph, R. B. Ruggeri, *Science* **260**, 1307 (1993).
- A. J. Robinson and P. B. Wyatt, *Tetrahedron* **49**, 11329 (1993).
- F. E. McDonald and S. J. Danishefsky, *J. Org. Chem.* **57**, 7001 (1992).
- C. Unverzagt and H. Kunz, *Bioorg. Med. Chem.* **3**, 197 (1993).
- M. Schuster, P. Wang, J. C. Paulson, C.-H. Wong, *J. Am. Chem. Soc.* **116**, 1135 (1994).
- S. J. Danishefsky, *Chemtracts Org. Chem.* **2**, 273 (1989); D. B. Berkowitz, S. J. Danishefsky, G. K. Schulte, *J. Am. Chem. Soc.* **114**, 4518 (1992).
- The polymer-bound disaccharide **18** was prepared from polymer-bound galactal carbonate and 3,4-galactal carbonate according to a procedure described in (17).
- R. U. Lemieux and S. Lavine, *Can. J. Chem.* **42**, 1473 (1964).
- S. Matsuura, C.-H. Niu, J. N. Cohen, *Chem. Commun.* **1976**, 451 (1976).
- H. Kunz and J. März, *Synlett* **1992**, 591 (1992).
- D. A. Jones Jr., *Tetrahedron Lett.* **33**, 2853 (1977).
- H. Paulsen, M. Schultz, D. Klamann, B. Waller, M. Paal, *Liebigs. Ann. Chem.* **1985**, 2028 (1985).
- This research was supported by NIH grant AI 16943. We also acknowledge a National Science and Engineering Research Council (Canada) postdoctoral fellowship to J.Y.R. and a NIH postdoctoral training grant T32 CA62948 to X.B.

9 February 1995; accepted 19 May 1995

Inhibitor-Enhanced Electron Transfer: Copper Cytochrome c as a Redox-Inert Probe of Ternary Complexes

Jian S. Zhou, Judith M. Nocek, Michael L. DeVan, Brian M. Hoffman*

Copper-substituted cytochrome c (CuCc) has been used as a structurally faithful, redox-inert inhibitor to probe the mechanism of electron transfer (ET) between Cc molecules and cytochrome c peroxidase (CcP). This inhibitor enhances photoinduced ET quenching of the triplet excited state of a zinc-substituted protein (ZnCcP or ZnCc) by its iron(III) partner (Fe³⁺Cc or Fe³⁺CcP). These results show that CcP and Cc form a ternary complex in which one Cc molecule binds tightly at a surface domain of CcP having low ET reactivity, whereas the second Cc molecule binds weakly to the 1:1 complex at a second domain with markedly greater (~10³) reactivity. These results also rule out the possibility that Cc bound at the second domain cooperatively enhances ET to Cc at the first domain. The multiphasic kinetics observed for the photoproduced ET intermediate do not reflect electron self-exchange between two Cc molecules within the ternary complex.

Respiration and metabolism depend on the sequential transfer of electrons from one protein to another (1), and this ET in turn

Department of Chemistry, Northwestern University, 2145 Sheridan Road, Evanston, IL 60208, USA.

*To whom correspondence should be addressed.

depends on the recognition and docking, as well as the reaction, of the ET partners (2). Issues of binding specificity and of reactivity in protein-protein reactions are analogous to those in enzyme-substrate reactions, but studies of ET complexes have lacked paral-